

Food and Drug Administration, HHS

§ 640.34

(2) Plasma may be obtained from a unit of Whole Blood collected by another licensed establishment.

[42 FR 59878, Nov. 22, 1977; 48 FR 13026, Mar. 29, 1983, as amended at 50 FR 4139, Jan. 29, 1985]

§ 640.31 Suitability of donors.

(a) Whole blood donors shall meet the criteria for donor suitability prescribed in § 640.3.

(b) Plasmapheresis donors shall meet the criteria for donor suitability prescribed in § 640.63, excluding the phrase “other than malaria” in paragraph (c)(9) of that section. Informed consent shall be required as prescribed in § 640.61.

[42 FR 59878, Nov. 22, 1977, as amended at 64 FR 45372, Aug. 19, 1999]

§ 640.32 Collection of source material.

(a) Whole blood shall be collected, transported, and stored as prescribed in § 640.4. When whole blood is intended for Plasma, Fresh Frozen Plasma, and Liquid Plasma, it shall be maintained at a temperature between 1 and 6 °C until the plasma is removed. Whole blood intended for Platelet Rich Plasma, shall be maintained as prescribed in § 640.24 until the plasma is removed. The red blood cells shall be placed in storage at a temperature between 1 and 6 °C immediately after the plasma is separated.

(b) Plasma obtained by plasmapheresis shall be collected as prescribed in §§ 640.62, 640.64 (except that paragraph (c)(3) of § 640.64 shall not apply), and § 640.65.

[42 FR 59878, Nov. 22, 1977, as amended at 45 FR 27927, Apr. 25, 1980; 50 FR 4139, Jan. 29, 1985; 64 FR 45372, Aug. 19, 1999]

§ 640.33 Testing the blood.

(a) Blood from which plasma is separated shall be tested as prescribed in § 610.40 of this chapter and § 640.5 (a), (b), and (c).

(b) Manufacturers of Plasma collected by plasmapheresis shall have testing and recordkeeping responsibilities

equivalent to those prescribed in §§ 640.71 and 640.72.

[42 FR 59878, Nov. 22, 1977, as amended at 44 FR 17658, Mar. 23, 1979; 50 FR 4139, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988; 66 FR 31165, June 11, 2001]

§ 640.34 Processing.

(a) *Plasma*. Plasma shall be separated from the red blood cells and shall be stored at -18 °C or colder within 6 hours after transfer to the final container or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system unless the product is to be stored as Liquid Plasma.

(b) *Fresh Frozen Plasma*. Fresh frozen plasma shall be prepared from blood collected by a single uninterrupted venipuncture with minimal damage to and minimal manipulation of the donor's tissue. The plasma shall be separated from the red blood cells, and placed in a freezer within 8 hours or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system, and stored at -18 °C or colder.

(c) *Liquid Plasma*. Liquid Plasma shall be separated from the red blood cells and shall be stored at a temperature of 1 to 6 °C within 4 hours after filling the final container or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system.

(d) *Platelet Rich Plasma*. Platelet rich plasma shall be prepared from blood collected by a single uninterrupted venipuncture with minimal damage to and manipulation of the donor's tissue. The plasma shall be separated from the red blood cells by centrifugation within 4 hours after completion of the phlebotomy or within the timeframe specified in the directions for use for the blood collecting, processing, and storage system. The time and speed of the centrifugation shall have been shown to produce a product with at least 250,000 platelets per microliter. The plasma shall be stored at a temperature between 20 and 24 °C immediately after filling the final container. A gentle and continuous agitation of the